arXiv:1211.4735v2 [cond-mat.str-el] 28 Nov 2012

The phase coherence and pseudogap at the excitonic phase transition in a three-dimensional extended Falicov-Kimball model

V. A. Apinyan and T. K. Kopeć
Institute for Low Temperature and Structure Research,
Polish Academy of Sciences
PO. Box 1410, 50-950 Wroclaw 2, Poland

(Dated: November 29, 2012)

Considering the phase coherence mechanism between the conduction band electrons and valence band holes we derive the excitonic phase stiffness for the three-dimensional strongly correlated electron system. It is shown that the phase coherence between the conduction and valence bands is important for the excitonic condensation at the very low temperatures. We prove that the Bose-superfluid excitonic transition critical lines driven by the phase stiffness in the strong and intermediate coupling regimes differ considerably from the excitonic pair formation temperature transition lines. The true condensate transition temperature appears to be much smaller than the exciton pair formation transition temperature. We show that a BEC to BCS type crossover could be expected in our model. In the strong interaction case the system is driven into the insulator state with a BEC condensation at the very low temperatures, while in the small interaction region the condensate order is governed by the pair formation and in this case the system is evidently in the BCS-like regime.

PACS numbers: Valid PACS appear here

I. INTRODUCTION

The Coulomb interaction between the conduction band electrons and the valence band holes causes in the solid state materials the formation of the new bound states of these two quasiparticles called the excitons.[1] The low-density system of excitons behaves like a weakly nonideal Bose-gas.[2] These new formations lead to the various interesting physical phenomena in solid state materials and they are the subjects of the intensive experimental and theoretical researches.[3–9] The ground state of the strongly interacting solid state system is unstable against the formation of the excitonic pairs and at the very low temperatures the new physical phases of the matter could develop.[1] Particularly, in the scenario of the semiconductor-metal phase transition a new phase develops approaching to the transition from the semiconductor side.[8] This state is called as the "excitonic insulator" (EI) state and is characterized by the strong coherence between the conduction band electrons and valence band holes. In the EI state a gap for the charged quasiparticle excitations is always present.[1] It has been shown[10] that the transition into the EI state is a second-order phase transition. Series of the recent experimental investigations[3,4] in $\mathrm{TmSe_{0.45}Te_{0.55}}$ suggested the existence of the EI state in that material. Another example of the material with a well defined EI state is the quasi-one-dimensional $\mathrm{Ta_2NiSe_5}$ with highly polarizable Se. The angle-resolved photoemission spectra on these compounds[11] demonstrate that the ground state therein is an excitonic insulator. In general the solid state system in the EI state is considered as an electron-hole pair condensate[1] at sufficiently low temperatures. In the small interaction region or when approaching to the EI state from the metal part of the transition spectrum, the system is in the Bardeen-Cooper-Schrieffer[12](BCS) superconducting condensate like regime with weakly bound electron-hole pairs and approaching from the semiconductor side the system shows typical Bose-Einstein condensation[13](BEC) behavior with the tightly bound excitons. Thus in the systems with the EI phase instability a BCS-BEC crossover[14] could be expected. This type of crossover mechanism is discussed in Ref. 15, where the authors find a transition from the BEC of excitons to a BCS state of electron-hole pairs. Using the screened Coulomb potential they derive the equations of quantum condensation using the real time Green function technique for a partially equilibrium system. In the context of the EI state, the three-dimensional spinless Falicov-Kimball model with the dispersive f-orbital electrons at half-filling has been analyzed recently.[16] Particularly, from the behavior of the partial f and c-electron densities of states, the authors find a crossover from a superconducting BCS-like condensate to a Bose-Einstein type excitonic condensate. A finite temperature transition critical line is established when the Hartree gap is non zero. The spontaneous symmetry breaking for the EI state and the BCS-BEC like crossover for the two-dimensional extended Falicov-Kimball model (EFKM) is discussed also in Ref. 17. The authors study the EI state using the variational cluster approximation (VCA) and the Hartree-Fock (HF) approximation. It is shown that the coherence length of the excitonic pair spread wide for several hundred lattice spacing. This indicates that the excitonic insulator state typifies either a BCS-like weakly-bound condensate of electron-hole pairs or a Bose-Einstein condensate of preformed excitons. The nature of the electron-hole bound states is analyzed in Ref. 18. Especially, at the semimetal-semiconductor transition the electron-hole pair condensation is found and the precursor effects for the electron-hole pair susceptibility is considered.

At the temperatures below the temperature of the quantum degeneracy the excitonic quasiparticles obeying the Bose-statistics can spontaneously form the coherent states, which is the main characteristic of condensation in momentum space. At this level of treatment, to explore the quantum physics of the excitons and in contrast with the mentioned theoretical works above,[15–18] we derive the EI state by showing its direct relation to the phase coherence between the conduction band electrons and valence band holes. We show that this coherence is leading furthermore to the exciton Bose-condensation at the sufficiently low temperatures. Following the original papers,[19,20] our model contains two type of fermions. The valence band f-electrons with the orbital energy ϵ_f couple to the itinerant c-electrons with the orbital energy ϵ_c supposing that there is a mechanism for the f-c coherence. In the same spirit of the works,[16,17] we discuss the EFKM in the context of the path integral formalism and we derive the EI state solving the system of self-consistent set of equations. Basing on the form of the effective phase stiffness action derived in the model, we prove that the phase coherence between the c-orbital electrons and f-band holes is essential along the whole pair formation EI line. The important issue of the theory is the derivation of the excitonic condensate order. The temperature transition critical lines are derived employing the effective quantum rotor model. We show that temperature transition critical lines obtained for the Bose-superfluid condensation of excitons in the three-dimensional system of excitons are considerably different from the excitonic pair formation temperature transition lines. Hence, we show that pair formation transition and the excitonic condensate transition are two different lines which are in coincidence only in the very small Coulomb interaction region, where the system is driven to the BCS-like state. Our theory provides an alternative mechanism for the BEC-BCS crossover discussed earlier.

The present article is organized as follows. In the Section II we provide the Hamiltonian of the model EFKM, then in the Section III we introduce the new decoupling potentials and we handle with the four fermion interaction term in the Hamiltonian. In Section IV we derive the excitonic pairing gap parameter, single-particle excitation energies and the average particle concentrations for each band. In Section V we derive the phase stiffness parameter and the self-consistent equation for the excitonic condensate transition temperature in EFKM and we find the complete quantum critical solution of the EFKM model.

II. THE HAMILTONIAN

We consider the Hamiltonian of the extended spinless Falicov-Kimball model in which the electrons with one spin direction are localized, while those with the other spin direction are moving

$$\begin{aligned}\mathcal{H} &= -t\sum_{\langle\mathbf{rr}'\rangle}\left[\bar{c}(\mathbf{r})c(\mathbf{r}')+h.c.\right]-(\mu-\epsilon_c)\sum_{\mathbf{r}}n_c(\mathbf{r})\\ &-\tilde{t}\sum_{\langle\mathbf{rr}'\rangle}\left[\bar{f}(\mathbf{r})f(\mathbf{r}')+h.c.\right]-(\mu-\epsilon_f)\sum_{\mathbf{r}}n_f(\mathbf{r})\\ &+U\sum_{\mathbf{r}}n_c(\mathbf{r})n_f(\mathbf{r}),\end{aligned}\tag{1}$$

where $\langle\mathbf{rr}'\rangle$ runs over the nearest neighbor sites, $\bar{c}(\mathbf{r})$ ($c(\mathbf{r})$) are the creation (annihilation) operators of the electron of the c-orbitals at the site with the position $\mathbf{r}$, t is the hopping integral for the c-electrons and ϵ_c is the on-site energy level. Similarly, $\bar{f}(\mathbf{r})$ ($f(\mathbf{r})$) are the creation (annihilation) operators of the f-orbital electrons and $\tilde{t}$ is the hopping integral for the f-electrons. Furthermore, ϵ_f is the on-site energy level of the f-orbital. The chemical potential μ entering in Eq.(1) is defined for our grand canonical system. We suppose that the chemical potentials of both orbitals are the same as in the work in Ref. 17. Parameter U, entering in the last term of the Hamiltonian, is the Coulomb repulsion between two types of electrons. $n_c(\mathbf{r})$ and $n_f(\mathbf{r})$ are the c- and f-electron density operators and they are defined as usual by the relation $n_x(\mathbf{r}) = \bar{x}(\mathbf{r})x(\mathbf{r})$ with $x = c, f$ for the c and f-orbital electrons respectively. We will use throughout the paper the units $\hbar = k_B = 1$. We will use $t = 1$ as the measure of the energy. We consider also the following values for the band parameters $\epsilon_c = 0$ and $\epsilon_f = -1$ also in accordance with the Ref. 17. With this consideration the c- electrons are itinerant and the f-electrons are quasilocalized on the atomic sites. The equilibrium value of the chemical potential will be determined from the half-filling condition (the grand canonical approach) of the average particle density functions, i.e. $n_c = 1-n_f$, where $n_x \equiv \langle n_x(\mathbf{r})\rangle$ is the average particle density.

III. DECOUPLING PROCEDURE AND GAUGE FIELDS

We will decouple the four fermionic term in the Hamiltonian Eq.(1) within a path integral formalism. For this, we rewrite first the density product in the last term in Eq.(1) in the equivalent form

$$n_c(\mathbf{r})n_f(\mathbf{r}) = \frac{n^2(\mathbf{r})}{2}-\frac{\tilde{n}^2(\mathbf{r})}{2},\tag{2}$$

where we introduced the short-hand notations

$$n(\mathbf{r}) = n_c(\mathbf{r})+n_f(\mathbf{r}),\tag{3}$$

$$\tilde{n}(\mathbf{r}) = n_c(\mathbf{r})-n_f(\mathbf{r}).\tag{4}$$

With the new notations we can rewrite the Hamiltonian Eq.(1) of the system as

$$\begin{aligned} \mathcal{H} = &-t\sum_{\langle \mathbf{r}\mathbf{r}'\rangle}\left[\bar{c}(\mathbf{r})c(\mathbf{r}') + h.c.\right] - \bar{\mu}\sum_{\mathbf{r}} n(\mathbf{r}) \\ &-\tilde{t}\sum_{\langle \mathbf{r}\mathbf{r}'\rangle}\left[\bar{f}(\mathbf{r})f(\mathbf{r}') + h.c.\right] + \frac{\epsilon_c-\epsilon_f}{2}\sum_{\mathbf{r}}\tilde{n}(\mathbf{r}) \\ &+U\sum_{\mathbf{r}}\frac{1}{2}\left[n^2(\mathbf{r}) - \tilde{n}^2(\mathbf{r})\right]. \end{aligned} \tag{5}$$

We put here $\bar{\mu} = \mu - \bar{\epsilon}$ and $\bar{\epsilon} = (\epsilon_c + \epsilon_f)/2$ is the average band energy parameter.

A. Functional integral formulation

Dealing with the fermions, it is convenient to introduce the Grassmann variables $c(\mathbf{r}\tau)$ and $f(\mathbf{r}\tau)$ at each site $\mathbf{r}$ and imaginary time τ varying in the interval $0 \leq \tau \leq \beta$, where $\beta = 1/k_B T$ (with T being the thermodynamic temperature). The variables $c(\mathbf{r}\tau)$ and $f(\mathbf{r}\tau)$ satisfy the anti-periodic boundary conditions $x(\mathbf{r}\tau) = -x(\mathbf{r}\tau + \beta)$. The partition function of the system of fermions written as a functional integral over the Grassmann field is

$$\mathcal{Z} = \int\left[\mathcal{D}\bar{c}\mathcal{D}c\right]\int\left[\mathcal{D}\bar{f}\mathcal{D}f\right] e^{-\mathcal{S}[\bar{c},c,\bar{f},f]}, \tag{6}$$

where the action in the exponential is given in the path integral formulation as

$$\mathcal{S}[\bar{c},c,\bar{f},f] = \mathcal{S}_B[\bar{c},c] + \mathcal{S}_B[\bar{f},f] + \int_0^\beta d\tau \mathcal{H}(\tau). \tag{7}$$

Here $\mathcal{S}_B[\bar{c},c]$ and $\mathcal{S}_B[\bar{f},f]$ are the fermionic Berry terms for the c- and f-electrons respectively. They are defined as

$$\mathcal{S}_B[\bar{x},x] = \sum_{\mathbf{r}}\int_0^\beta d\tau \bar{x}(\mathbf{r}\tau)\frac{\partial}{\partial\tau}x(\mathbf{r}\tau). \tag{8}$$

Then we decouple the quadratic density terms in Eq.(5) using the Hubbard-Stratanovich transformation and by introducing the new decoupling variables $\mathrm{V}(\mathbf{r}\tau)$ and $\varrho(\mathbf{r}\tau)$ conjugated to the electron density operators $n(\mathbf{r}\tau)$ and $\tilde{n}(\mathbf{r}\tau)$. For the quadratic term in the exponential of the partition function proportional to $n^2(\mathbf{r}\tau)$ we have

$$\begin{aligned} &\exp\left(-\frac{U}{2}\sum_{\mathbf{r}}\int_0^\beta d\tau n^2(\mathbf{r}\tau)\right) \\ &= \int\left[\mathcal{D}\mathrm{V}\right] e^{-\sum_{\mathbf{r}}\int_0^\beta d\tau\left[\frac{\mathrm{V}^2(\mathbf{r}\tau)}{2U} - i\mathrm{V}(\mathbf{r}\tau)n(\mathbf{r}\tau)\right]}. \end{aligned} \tag{9}$$

The decoupling of the quadratic term proportional to $\tilde{n}^2(\mathbf{r}\tau)$ in the exponential of the partition function in Eq.(6) is also straightforward. We have

$$\begin{aligned} &\exp\left(\sum_{\mathbf{r}}\int_0^\beta d\tau\frac{U}{2}\tilde{n}^2(\mathbf{r}\tau)\right) \\ &= \int\left[\mathcal{D}\varrho\right] e^{-\sum_{\mathbf{r}}\int_0^\beta d\tau\left[\frac{\varrho^2(\mathbf{r}\tau)}{2U} - \varrho(\mathbf{r}\tau)\tilde{n}(\mathbf{r}\tau)\right]}. \end{aligned} \tag{10}$$

Shifting the integration variables in Eq.(10) we will have

$$\int\left[\mathcal{D}\varrho\right] e^{\sum_{\mathbf{r}}\int_0^\beta d\tau\left\{-\frac{\varrho^2(\mathbf{r}\tau)}{2U} - \varrho(\mathbf{r}\tau)\left[\tilde{n}(\mathbf{r}\tau) - \frac{\epsilon_c-\epsilon_f}{2U}\right]\right\}}. \tag{11}$$

The saddle-point evaluation of the obtained integral gives for ϱ

$$\varrho_0 = U\tilde{n} - \frac{\epsilon_c - \epsilon_f}{2}, \tag{12}$$

where $\tilde{n} = \langle\tilde{n}(\mathbf{r}\tau)\rangle$. As the result of the decoupling we will have the following "Zeeman-like" contribution to the partition function

$$\exp\left(-\sum_{\mathbf{r}}\int_0^\beta d\tau\tilde{n}(\mathbf{r}\tau)\mu_\varrho\right) \tag{13}$$

with the effective chemical potential $\mu_\varrho = \frac{\epsilon_c-\epsilon_f}{2} - U\tilde{n}$. After combining the exponential in Eq.(9) with the effective $\bar{\mu}$ chemical potential term in Eq.(5) we can decompose the variables $\mathrm{V}(\mathbf{r}\tau)$ into a static and the periodic parts as

$$\mathrm{V}(\mathbf{r}\tau) = \mathrm{V}_0(\mathbf{r}) + \tilde{\mathrm{V}}(\mathbf{r}\tau). \tag{14}$$

The integration over $\mathrm{V}(\mathbf{r}\tau)$ variables now becomes the integration over the scalar static variables $\mathrm{V}_0(\mathbf{r})$ and the integration over the periodic fields $\tilde{\mathrm{V}}(\mathbf{r}\tau)$

$$\int\left[\mathcal{D}\mathrm{V}\right]... = \int\left[\mathcal{D}\mathrm{V}_0\right]\int\left[\mathcal{D}\tilde{\mathrm{V}}\right]... \tag{15}$$

For the periodic part in Eq.(14) we define the emergent Faraday-type relation introducing the U(1) topological phase field $\varphi(\mathbf{r}\tau)$

$$\tilde{\mathrm{V}}(\mathbf{r}\tau) = \frac{\partial\varphi(\mathbf{r}\tau)}{\partial\tau} \equiv \dot{\varphi}(\mathbf{r}\tau). \tag{16}$$

Thus, for the dynamic part, we transform the integration over the gauge variables $\tilde{\mathrm{V}}(\mathbf{r}\tau)$ into the integration over the generic phase variables $\varphi(\mathbf{r}\tau)$ in accordance with the relation in Eq.(16)

$$\int\left[\mathcal{D}\tilde{\mathrm{V}}\right]... \to \int\left[\mathcal{D}\varphi\right]... \tag{17}$$

In performing the integration over the phase field one should take into account that the field configurations satisfy the condition

$$\varphi(\mathbf{r}\beta) - \varphi(\mathbf{r}0) = 2\pi m(\mathbf{r}), \tag{18}$$

where $m(\mathbf{r}) = 0, \pm1, \pm2, ...$ are the U(1) topological winding numbers, which are defining a set of the equivalent topological sectors. Thus the variables $\varphi(\mathbf{r}\tau)$ only with discrete configurations are matter. The proper integration measure in Eq.(17) over φ variables is a multiply-connected topological domain and

$$\int [\mathcal{D}\varphi] ... \equiv \sum_{[m(\mathbf{r})]} \int_0^{2\pi} \prod_{\mathbf{r}} d\varphi_0(\mathbf{r}) \times \\ \times \int_{\varphi(\mathbf{r}0)=\varphi(\mathbf{r})}^{\varphi(\mathbf{r}\beta)=\varphi(\mathbf{r})+2\pi m(\mathbf{r})} \prod_{\mathbf{r}} d\varphi(\mathbf{r}\tau) ..., \tag{19}$$

where in each topological sector the integration goes over the gauge potentials with the winding number equal to $m(r)$. The periodic part, on the other hand, can be decomposed into the Fourier series as

$$\tilde{\mathrm{V}}(\mathbf{r}\tau) = \frac{1}{\beta} \sum_{n=1}^{\infty} \left[\tilde{\mathrm{V}}(\mathbf{r}\omega_n) e^{i\omega_n \tau} + c.c. \right] \tag{20}$$

with $\omega_n = 2\pi n/\beta$ $(n = 0, \pm1, \pm2, ...)$ being the Bose-Matsubara frequencies. The decomposition of the charge field conforms with the basic $m = 0$ topological sector, because

$$\int_0^\beta d\tau \frac{\partial \varphi(\mathbf{r}\tau)}{\partial \tau} = \int_0^\beta d\tau \tilde{\mathrm{V}}(\mathbf{r}\tau) \equiv 0. \tag{21}$$

For the scalar static part $V_0(\mathbf{r})$ we have the following functional integral

$$\int [d\mathrm{V}_0]\, e^{\sum_{\mathbf{r}} \int_0^\beta d\tau \left\{ -\frac{\mathrm{V}_0^2(\mathbf{r})}{2U} + i\mathrm{V}_0(\mathbf{r})[n(\mathbf{r}\tau) - U\bar{\mu}] \right\}}. \tag{22}$$

The saddle-point value for the static part $\mathrm{V}_0(\mathbf{r})$ is given as $\mathrm{V}_0^{\mathrm{sp}} = iUn - i\bar{\mu}$, where n is the total average particle density $n = n_c + n_f$. And we have the contribution to the partition function in Eq.(6) in the form

$$\int [\mathcal{D}\varphi]\, e^{-\mathcal{S}[\varphi]}, \tag{23}$$

where the effective phase only action $\mathcal{S}[\varphi]$ is given as

$$\mathcal{S}[\varphi] = \sum_{\mathbf{r}} \int_0^\beta d\tau \left[\frac{\dot{\varphi}^2(\mathbf{r}\tau)}{2U} - \frac{\bar{\mu}}{iU} \dot{\varphi}(\mathbf{r}\tau) - i\dot{\varphi}(\mathbf{r}\tau) n(\mathbf{r}\tau) \right]. \tag{24}$$

Combining the exponentials in Eq.(13) and in Eq.(22) (after saddle-point evaluations) we will have our decoupling result in the form

$$\exp\left(-\sum_{\mathbf{r}} \int_0^\beta d\tau \left[\tilde{\mu} n(\mathbf{r}\tau) + \mu_\varrho \tilde{n}(\mathbf{r}\tau) \right] \right) \int [\mathcal{D}\varphi]\, e^{-\mathcal{S}[\varphi]}, \tag{25}$$

where the effective chemical potential $\tilde{\mu}$ is defined as $\tilde{\mu} = Un - \bar{\mu}$. The partition function of the system after decoupling procedures will be

$$\mathcal{Z} = \int [\mathcal{D}\bar{c}\mathcal{D}c] \left[\mathcal{D}\bar{f}\mathcal{D}f\right] [\mathcal{D}\varphi]\, e^{-\mathcal{S}[\bar{c},c,\bar{f},f,\varphi]}, \tag{26}$$

where the action $\mathcal{S}[\bar{c}, c, \bar{f}, f, \varphi]$ in the exponential is

$$\begin{aligned} \mathcal{S}[\bar{c}, c, \bar{f}, f, \varphi] &= \mathcal{S}[\varphi] + \mathcal{S}_B[\bar{c}, c] + \mathcal{S}_B[\bar{f}, f] \\ &- t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \int_0^\beta d\tau \left[\bar{c}(\mathbf{r}\tau) c(\mathbf{r}'\tau) + h.c. \right] \\ &- \tilde{t} \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \int_0^\beta d\tau \left[\bar{f}(\mathbf{r}\tau) f(\mathbf{r}'\tau) + h.c. \right] \\ &+ \sum_{\mathbf{r}} \int_0^\beta d\tau \left[\tilde{\mu} n(\mathbf{r}\tau) + \mu_\varrho \tilde{n}(\mathbf{r}\tau) \right]. \end{aligned} \tag{27}$$

We see that a strongly fluctuating imaginary term appear in the expression of the phase action in Eq.(24)). In order to eliminate this term we will apply the U(1) gauge transformations to the fermionic variables $c(\mathbf{r}\tau)$ and $f(\mathbf{r}\tau)$.

B. The U(1) gauge transformation

We perform the local gauge transformation to the new fermionic Grassmann variables $a(\mathbf{r}\tau)$ and $b(\mathbf{r}\tau)$ in order to eliminate the last imaginary term appearing in the expression of the phase action in Eq.(24). Doing so, we cast the strongly correlated fermionic system into a weakly interacting one, which is submerged into the bath of strongly fluctuating gauge potentials (whose interactions are governed by a high energy scales of U). For the c-orbital electrons the U(1) transformation is

$$\begin{bmatrix} c(\mathbf{r}\tau) \\ \bar{c}(\mathbf{r}\tau) \end{bmatrix} = \begin{bmatrix} e^{i\varphi(\mathbf{r}\tau)} & 0 \\ 0 & e^{-i\varphi(\mathbf{r}\tau)} \end{bmatrix} \begin{bmatrix} a(\mathbf{r}\tau) \\ \bar{a}(\mathbf{r}\tau) \end{bmatrix}, \tag{28}$$

where we introduced the new bosonic phase variables φ. For the f-orbital electrons the similar transformation is

$$\begin{bmatrix} f(\mathbf{r}\tau) \\ \bar{f}(\mathbf{r}\tau) \end{bmatrix} = \begin{bmatrix} e^{i\varphi(\mathbf{r}\tau)} & 0 \\ 0 & e^{-i\varphi(\mathbf{r}\tau)} \end{bmatrix} \begin{bmatrix} b(\mathbf{r}\tau) \\ \bar{b}(\mathbf{r}\tau) \end{bmatrix}. \tag{29}$$

After those transformations the electronic action of the system in Eq.(27) takes the following form

$$\begin{aligned} \mathcal{S}[\bar{a}, a, \bar{b}, b, \varphi] &= \mathcal{S}_0[\varphi] + \mathcal{S}_B\left[\bar{a}, a\right] + \mathcal{S}_B\left[\bar{b}, b\right] \\ &- t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \int_0^\beta d\tau \left[\bar{a}(\mathbf{r}\tau) a(\mathbf{r}'\tau) e^{-i\left[\varphi(\mathbf{r}\tau) - \varphi(\mathbf{r}'\tau)\right]} + h.c. \right] \\ &- \tilde{t} \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \int_0^\beta d\tau \left[\bar{b}(\mathbf{r}\tau) b(\mathbf{r}'\tau) e^{-i\left[\varphi(\mathbf{r}\tau) - \varphi(\mathbf{r}'\tau)\right]} + h.c. \right] \\ &+ \sum_{\mathbf{r}} \int_0^\beta d\tau \left[\tilde{\mu} n(\mathbf{r}\tau) + \mu_\varrho \tilde{n}(\mathbf{r}\tau) \right]. \end{aligned} \tag{30}$$

The phase action $\mathcal{S}_0[\varphi]$ in Eq.(30) is derived as

$$\mathcal{S}_0[\varphi]=\sum_{\mathbf{r}}\int_0^\beta d\tau\left[\frac{\dot{\varphi}^2(\mathbf{r}\tau)}{2U}-\frac{\bar{\mu}}{iU}\dot{\varphi}(\mathbf{r}\tau)\right]. \qquad (31)$$

The partition function of the system in new variables is

$$\mathcal{Z}=\int\left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right]\left[\mathcal{D}\varphi\right]e^{-\mathcal{S}[\bar{a},a,\bar{b},b,\varphi]}. \qquad (32)$$

This form of the partition function is very general and furthermore we will use it for deriving the effective actions for the fermions and for the phase sector.

IV. RESULTS OF CALCULATIONS

A. Effective fermionic action and propagators

We start with the phase averaged expression of the action in Eq.(7) after the U(1) transformations given by the Eqs.(28) and (29). Applying the Hartree-Fock (HF) like approximation (see Ref. 23) to the four-fermion term in the Hamiltonian of Eq.(1), the phase averaged expression of the total action can be derived. We present here the Fourier space evaluations (the derivation in the real space is very similar). The Fourier transformation of the fermionic variables $a(\mathbf{r}\tau)$ and $b(\mathbf{r}\tau)$ is given by

$$x(\mathbf{r}\tau)=\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}x_{\mathbf{k}}(\nu_n)e^{i(\mathbf{kr}-\nu_n\tau)} \qquad (33)$$

with $x=a,b$ for the a and b type electrons. N is the total number of $\mathbf{k}$-points in the first Brillouin zone (FBZ) and $\nu_n=\pi(2n+1)/\beta$ are the Fermi-Matsubara frequencies with $n=0,\pm1,\pm2,\ldots$. We introduce the local excitonic gap parameter Δ as in the Ref. 17

$$\Delta=U\left\langle\bar{a}(\mathbf{r}\tau)b(\mathbf{r}\tau)\right\rangle. \qquad (34)$$

Then, the HF decoupling of the interaction term in the Hamiltonian in Eq.(1) is

$$n_a(\mathbf{r}\tau)n_b(\mathbf{r}\tau)\approx\left\langle n_a(\mathbf{r}\tau)\right\rangle n_b(\mathbf{r}\tau)+\left\langle n_b(\mathbf{r}\tau)\right\rangle n_a(\mathbf{r}\tau)$$
$$-\frac{1}{U}\bar{\Delta}(\mathbf{0},0)\bar{a}(\mathbf{r}\tau)b(\mathbf{r}\tau)-\frac{1}{U}\Delta(\mathbf{0},0)\bar{b}(\mathbf{r}\tau)a(\mathbf{r}\tau), \qquad (35)$$

where $n_a(\mathbf{r}\tau)$ and $n_b(\mathbf{r}\tau)$ are the electron density operators after the U(1) gauge transformation. Due to the symmetry of the U(1) transformation they are practically the same operators as $n_c(\mathbf{r}\tau)$ and $n_f(\mathbf{r}\tau)$, because of the relations $n_a(\mathbf{r}\tau)\equiv n_c(\mathbf{r}\tau)$ and $n_b(\mathbf{r}\tau)\equiv n_f(\mathbf{r}\tau)$. Then action of the system takes the form

$$\begin{aligned}\mathcal{S}\left[\bar{a},a,\bar{b},b\right]=&-\frac{\bar{\Delta}}{\beta N}\sum_{\mathbf{k}\nu_n}\bar{a}_{\mathbf{k}}(\nu_n)b_{\mathbf{k}}(\nu_n)\\&-\frac{\Delta}{\beta N}\sum_{\mathbf{k}\nu_n}\bar{b}_{\mathbf{k}}(\nu_n)a_{\mathbf{k}}(\nu_n)\\&+\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}\bar{a}_{\mathbf{k}}(\nu_n)\left(\mu^a_{\rm eff}-i\nu_n-t_{\mathbf{k}}\right)a_{\mathbf{k}}(\nu_n)\\&+\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}\bar{b}_{\mathbf{k}}(\nu_n)\left(\mu^b_{\rm eff}-i\nu_n-\tilde{t}_{\mathbf{k}}\right)b_{\mathbf{k}}(\nu_n),\end{aligned} \qquad (36)$$

where $\Delta\equiv\Delta(\mathbf{0},0)$ is the local excitonic gap parameter and the effective chemical potentials $\mu^a_{\rm eff}$ and $\mu^b_{\rm eff}$ are defined as

$$\mu^a_{\rm eff}=\epsilon_a-\mu+Un_b+i\left\langle\dot{\varphi}(\mathbf{r}\tau)\right\rangle, \qquad (37)$$
$$\mu^b_{\rm eff}=\epsilon_b-\mu+Un_a+i\left\langle\dot{\varphi}(\mathbf{r}\tau)\right\rangle. \qquad (38)$$

n_a and n_b in the definitions of Eq.(37) and Eq.(38) are the average values for the respective density operators and $n_x=\langle n_x(\mathbf{r}\tau)\rangle$ with $x=a,b$. The renormalized hopping amplitudes $t_{\mathbf{k}}$ and $\tilde{t}_{\mathbf{k}}$ are given by the relations

$$t_{\mathbf{k}}=2tg\epsilon\left(\mathbf{k}\right), \qquad (39)$$
$$\tilde{t}_{\mathbf{k}}=2\tilde{t}g\epsilon\left(\mathbf{k}\right), \qquad (40)$$

where

$$\epsilon\left(\mathbf{k}\right)=\cos(d_xk_x)+\cos(d_yk_y)+\cos(d_zk_z) \qquad (41)$$

is the three-dimensional lattice dispersion relation with d_i, $i=x,y,z$, being the components of the lattice spacing vector $\mathbf{d}=\mathbf{r}-\mathbf{r}'$ with $\mathbf{r}$ and $\mathbf{r}'$ nearest neighbors sites. The bandwidth renormalization factor g is given as

$$g=\left\langle e^{-i[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau)]}\right\rangle\Big|_{|\mathbf{r}-\mathbf{r}'|=\mathbf{d}}. \qquad (42)$$

Introducing the vector space notations, we can rewrite the action in Eq.(36) in the form

$$\mathcal{S}\left[\bar{a},a,\bar{b},b\right]=\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}\left[\bar{a}_{\mathbf{k}}(\nu_n),\bar{b}_{\mathbf{k}}(\nu_n)\right]\mathcal{G}^{-1}(\mathbf{k}\nu_n)\times$$
$$\times\left[\begin{array}{c}a_{\mathbf{k}}(\nu_n)\\b_{\mathbf{k}}(\nu_n)\end{array}\right]. \qquad (43)$$

Here $\mathcal{G}^{-1}(\mathbf{k}\nu_n)$ is the inverse of the Green function matrix given as

$$\mathcal{G}^{-1}(\mathbf{k}\nu_n)=\left[\begin{array}{cc}\mathcal{E}^a_{\mathbf{k}}(\nu_n)&-\bar{\Delta}\\-\Delta&\mathcal{E}^b_{\mathbf{k}}(\nu_n)\end{array}\right], \qquad (44)$$

where single particle quasienergies $\mathcal{E}^a_{\mathbf{k}}(\nu_n)$ and $\mathcal{E}^b_{\mathbf{k}}(\nu_n)$ are given after Eq.(36) as

$$\mathcal{E}^a_{\mathbf{k}}(\nu_n)=\mu^a_{\rm eff}-i\nu_n-t_{\mathbf{k}}, \qquad (45)$$
$$\mathcal{E}^b_{\mathbf{k}}(\nu_n)=\mu^b_{\rm eff}-i\nu_n-\tilde{t}_{\mathbf{k}}. \qquad (46)$$

The explicit expressions for the normal and anomalous excitonic propagators can be derived from the effective action in Eq.(43). The normal propagator is defined as usual by (see Ref. 24)

$$\mathcal{G}^{xx}(\mathbf{r}\tau,\mathbf{r}'\tau') = -\left\langle x(\mathbf{r}\tau)\bar{x}(\mathbf{r}'\tau')\right\rangle \tag{47}$$

The anomalous or the excitonic propagator is defined in the theory as

$$\mathcal{G}^{ab}(\mathbf{r}\tau,\mathbf{r}'\tau') = \left\langle \bar{a}(\mathbf{r}\tau)b(\mathbf{r}'\tau')\right\rangle . \tag{48}$$

Then, using the standard functional derivation technique,[21] we have for the normal electron propagators

$$\begin{aligned}\mathcal{G}^{aa}(\mathbf{r}\tau,\mathbf{r}'\tau') &= -\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}\mathcal{E}^{b}_{\mathbf{k}}(\nu_n)\frac{e^{i\left[\mathbf{k}(\mathbf{r}-\mathbf{r}')-\nu_n(\tau-\tau')\right]}}{\mathcal{E}^{a}_{\mathbf{k}}(\nu_n)\mathcal{E}^{b}_{\mathbf{k}}(\nu_n)-|\Delta|^2},\\ \mathcal{G}^{bb}(\mathbf{r}\tau,\mathbf{r}'\tau') &= -\frac{1}{\beta N}\sum_{\mathbf{k}\nu_n}\mathcal{E}^{a}_{\mathbf{k}}(\nu_n)\frac{e^{i\left[\mathbf{k}(\mathbf{r}-\mathbf{r}')-\nu_n(\tau-\tau')\right]}}{\mathcal{E}^{a}_{\mathbf{k}}(\nu_n)\mathcal{E}^{b}_{\mathbf{k}}(\nu_n)-|\Delta|^2}.\end{aligned} \tag{49}$$

The anomalous propagators are derived as

$$\begin{aligned}\mathcal{G}^{ab}(\mathbf{r}\tau,\mathbf{r}'\tau') &= -\frac{\bar{\Delta}}{\beta N}\sum_{\mathbf{k}\nu_n}\frac{e^{-i\left[\mathbf{k}(\mathbf{r}-\mathbf{r}')-\nu_n(\tau-\tau')\right]}}{\mathcal{E}^{a}_{\mathbf{k}}(\nu_n)\mathcal{E}^{b}_{\mathbf{k}}(\nu_n)-|\Delta|^2},\\ \mathcal{G}^{ba}(\mathbf{r}\tau,\mathbf{r}'\tau') &= -\frac{\Delta}{\beta N}\sum_{\mathbf{k}\nu_n}\frac{e^{-i\left[\mathbf{k}(\mathbf{r}-\mathbf{r}')-\nu_n(\tau-\tau')\right]}}{\mathcal{E}^{a}_{\mathbf{k}}(\nu_n)\mathcal{E}^{b}_{\mathbf{k}}(\nu_n)-|\Delta|^2}.\end{aligned} \tag{50}$$

The local expressions of the normal and excitonic propagators will be used to find the system of self-consistent equations of the theory. The anomalous propagators of the form given in Eq.(50) will determine the phase stiffness parameter (see Appendix B).

B. The excitonic gap parameter

The general definition of the excitonic order parameter is terms of the initial c-band electron and f-band hole operators $\bar{c}(\mathbf{r}\tau)$ and $f(\mathbf{r}\tau)$ is

$$\Psi = U\lim_{\mathbf{r}\to\mathbf{r}'}\left\langle \bar{c}(\mathbf{r}\tau)f(\mathbf{r}'\tau)\right\rangle . \tag{51}$$

Or after the U(1) transformations we have

$$\Psi = U\lim_{\mathbf{r}\to\mathbf{r}'}\left\langle \bar{a}(\mathbf{r}\tau)b(\mathbf{r}'\tau)\right\rangle \left\langle e^{i\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau)\right]}\right\rangle . \tag{52}$$

Thus we represented the excitonic order parameter Ψ as a product of the U(1) gap parameter $U\left\langle \bar{a}(\mathbf{r}\tau)b(\mathbf{r}'\tau')\right\rangle$ with a phase coherence transition probability function

$$\Psi = \lim_{\mathbf{r}\to\mathbf{r}'}\Delta(\mathbf{r}\tau,\mathbf{r}'\tau)\mathcal{G}_{\varphi}(\mathbf{r}-\mathbf{r}'), \tag{53}$$

where we introduced the excitonic gap parameter $\Delta(\mathbf{r}\tau,\mathbf{r}'\tau)$ as

$$\Delta(\mathbf{r}\tau,\mathbf{r}'\tau) = U\left\langle \bar{a}(\mathbf{r}\tau)b(\mathbf{r}'\tau)\right\rangle \tag{54}$$

and the probability function $\mathcal{G}_{\varphi}(\mathbf{r}-\mathbf{r}')$ is, in fact, a phase-phase correlation function defined as

$$\mathcal{G}_{\varphi}(\mathbf{r}-\mathbf{r}') = \left\langle e^{i\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau)\right]}\right\rangle . \tag{55}$$

As we will see in the Section V, only a nonzero value of the gap parameter is not sufficient for the possible prediction of the excitonic condensation. To achieve this, also the phase variables must become stiff and coherent, which implies that $\mathcal{G}_{\varphi}$ function defined in Eq.(55) is not vanishing at least for the fundamental state of the system in which the Bose particles are condensed in the $\mathbf{k} = 0$ momentum state of the reciprocal lattice space. This is the quantum state of the system, where a macroscopic number of particles are situated in the same state of the reciprocal space. The Fourier transformation of the phase coherence factor $\mathcal{G}_{\varphi}(\mathbf{r}-\mathbf{r}')$ could be written as a sum of two components, one of which is describing the condensate transition probability and the other one is just a usual k-summation over the reciprocal wave vectors, defining the non-condensed part. Hence we can write

$$\mathcal{G}_{\varphi}(\mathbf{r}-\mathbf{r}') = \psi_0^2 + \frac{1}{N}\sum_{\mathbf{k}\neq 0}e^{i\mathbf{k}(\mathbf{r}-\mathbf{r}')}\mathcal{G}_{\varphi}(\mathbf{k}), \tag{56}$$

where

$$\psi_0^2 = \frac{1}{N}\mathcal{G}_{\varphi}(\mathbf{0}) \tag{57}$$

is the condensate transition probability with

$$\psi_0 = \left\langle e^{i\varphi(\mathbf{r}\tau)}\right\rangle . \tag{58}$$

The non-zero value of the average in Eq.(58) is necessary for the transition to the Bose-superfluid state. The fact of the appearance of this non-vanishing expectation value leads to a phase transition from a symmetric phase $\psi_0 = 0$ to an unsymmetric phase $\psi_0 \neq 0$. We can write now the excitonic order parameter as

$$\Psi = \Delta\left[\psi_0^2 + \frac{1}{N}\sum_{\mathbf{k}\neq 0}\mathcal{G}_{\varphi}(\mathbf{k})\right], \tag{59}$$

where Δ is the local excitonic gap parameter $\Delta = U\left\langle \bar{a}(\mathbf{r}\tau)b(\mathbf{r}\tau)\right\rangle \equiv \Delta(\mathbf{0},0)$. Ψ order parameter is consisting of the condensate order plus the on-condensate quasiparticle excitations. Indeed, the second term in Eq.(59) gives contribution to the excitonic pairing mechanism without the condensate order. At the very low temperatures, at which the excitonic condensation could happens, we can neglect the on-condensate part in Eq.(59) and we have only the condensate order

$$\Psi = \Delta\psi_0^2. \tag{60}$$

Indeed, the generic U(1) gauge field variables φ introduced in Eq.(16), being apart the phase flux tubes attached to the fermionic variables, are also the main mechanism for the Bose-superfluid state in the system. The exact prove of it is given in the Section V, where the excitonic condensation is demonstrated using a quantum rotor model analogue to the 3D spherical model.[21]

C. Pairing gap

We will show below how the excitonic gap parameter Δ could be calculated by solving a system of self-consistent equations for our system and, later on, in the Section V we will derive the condensate order and show how the phase stiffness mechanism is responsible for the temperature transition critical lines to the condensate order in the system. From the expression of the total action of the system in Eq.(43), we can derive the equations for the average electron densities n_a and n_b corresponding to the a- and b-orbitals respectively and also a self-consistent equation for the excitonic gap parameter Δ. Then we will have

$$\begin{aligned} n_a &= \mathcal{G}^{aa}(\mathbf{0},0), \\ n_b &= \mathcal{G}^{bb}(\mathbf{0},0), \\ \Delta &= U\mathcal{G}^{ab}(\mathbf{0},0). \end{aligned} \tag{61}$$

Using the analytical expressions for the propagators given in Eq.(49) and Eq.(50), we can rewrite the system in Eq.(61) in an equivalent explicit form. We have

$$\frac{1}{N}\sum_{\mathbf{k}}\left[n_F(E^{+}_{\mathbf{k}})+n_F(E^{-}_{\mathbf{k}})\right]=1, \tag{62}$$

$$\tilde{n}=\frac{1}{N}\sum_{\mathbf{k}}\xi_{\mathbf{k}}\frac{n_F(E^{+}_{\mathbf{k}})-n_F(E^{-}_{\mathbf{k}})}{\sqrt{\xi^2_{\mathbf{k}}+4\Delta^2}}, \tag{63}$$

$$\Delta=-\frac{U\Delta}{N}\sum_{\mathbf{k}}\frac{n_F(E^{+}_{\mathbf{k}})-n_F(E^{-}_{\mathbf{k}})}{\sqrt{\xi^2_{\mathbf{k}}+4\Delta^2}}, \tag{64}$$

where we used the fact that at half-filling $n=n_a+n_b=1$ and we defined $\tilde{n}=n_a-n_b$. n_F is the Fermi-Dirac distribution function $n_F(\epsilon)=1/(e^{\beta\epsilon}+1)$ and the energy parameters $E^{+}_{\mathbf{k}}$ and $E^{-}_{\mathbf{k}}$ are given by

$$E^{\pm}_{\mathbf{k}}=\frac{1}{2}\left(-A_{\mathbf{k}}\pm\sqrt{\xi^2_{\mathbf{k}}+4\Delta^2}\right), \tag{65}$$

where $A_{\mathbf{k}}=\tilde{t}_{\mathbf{k}}-\mu^{b}_{\mathrm{eff}}+t_{\mathbf{k}}-\mu^{a}_{\mathrm{eff}}$ and the quasiparticle dispersion is $\xi_{\mathbf{k}}=\tilde{t}_{\mathbf{k}}+\mu^{a}_{\mathrm{eff}}-t_{\mathbf{k}}-\mu^{b}_{\mathrm{eff}}$. The quantities n_a, n_b, Δ and μ can be determined by solving numerically the above equations in a self-consistent way. From the summations over the wave vectors in Eqs.(62), (63) and (64) we can pass to the integrations by introducing the appropriate density of states (DOS) for the three-dimensional lattice

$$\rho_{3D}(\zeta)=\frac{1}{N}\sum_{\mathbf{k}}\delta\left[\zeta-\sum_{\mathbf{d}}\cos(\mathbf{kd})\right]. \tag{66}$$

For the simple cubic lattice geometry the density of states is given as

$$\begin{aligned}\rho_{3D}(\zeta)=\frac{1}{\pi^3}\int_{\max(-1,-2-\zeta)}^{\min(1,2-\zeta)}dy\frac{\Theta\left(1-\frac{|\zeta|}{3}\right)}{\sqrt{1-y^2}}\times \\ \times\mathbf{K}\left[\sqrt{1-\left(\frac{y}{2}+\frac{\zeta}{2}\right)^2}\right],\end{aligned} \tag{67}$$

where $\Theta(x)$ is the Heaviside step function and $\mathbf{K}(x)$ is the elliptic function of the first kind.[25]The plot of the function $\rho_{3D}(x)$ is presented in the inset of Fig. 1, where the electronic density difference function $\tilde{n}$ is plotted as a function of the normalized Coulomb interaction parameter U for the excitonic pair formation region, i.e. when $\Delta\neq 0$. Two different values of the hopping parameter $\tilde{t}$ are considered and the solutions at $T=0$ K are presented. It is clear from Fig. 1 that in the strong coupling limit, i.e. for the higher values of the Coulomb interaction, the system is in the insulator regime, because at the upper boundary of the Coulomb interaction parameter the b-band is fully occupied ($n_b=1$) and the a-band is totally free ($n_a=0$). It is also clear from Fig. 1 that with increasing the hopping integral the system reaches the "excitonic insulator" phase at the higher values of the Coulomb interaction. This is due to the fact that the higher hopping amplitudes are destabilizing the insulator phase and the limit $n_a=0$ is achieved in the strong coupling case. In Fig. 2 the excitonic pairing gap parameter is plotted as a function of the Coulomb interaction energy in units of t and for different values of the hopping amplitude $\tilde{t}$ equal to $\tilde{t}=-0.3$ for the upper curve and equal to $\tilde{t}=-0.1$ for the lower curve. Decreasing (in absolute value) the hopping integral $\tilde{t}$, we observe that the correlation region becomes narrow, vanishing for the smaller values of U/t. The correlation amplitude also has been reduced. It is clear that in the three-dimensional case we have the long-range correlations between the electrons and holes. The excitonic gap is not vanishing for a rather large domain of the Coulomb interaction in difference with the results for the two-dimensional square lattice geometry given in Ref. 17. The authors in Ref. 17, analyzing the EFKM Hamiltonian using the Hartree-Fock approximation, derive the gap parameter for a narrow range of the Coulomb interaction strength U. The obtained values for the lower and upper bounds of the Coulomb interaction in Ref. 17 are about $(U_{c1},U_{c2})=(0.66,6.95)$, which differ considerably from our results especially for the large hoppings (see the upper curve on Fig. 2 for $\tilde{t}=-0.3$). In our case, the values for the lower and upper bounds of the normalized Coulomb interaction are given



FIG. 1: The average particle density difference function $\tilde{n}$ as a function of the dimensionless interaction parameter U/t for two different values of the hopping integral $\tilde{t}$ and at $T = 0$ K. In the inset of the figure the density of states (DOS) for the three-dimensional cubic lattice is presented.



FIG. 2: The excitonic gap parameter Δ as a function of the normalized Coulomb interaction energy U (in units of t) at $T = 0$ K for two different hopping parameters.

with the intervals $(U_{c1}, U_{c2}) = (0.7, 10.7)$ for $\tilde{t} = -0.3$ and $(U_{c1}, U_{c2}) = (0.8, 7.8)$ for the case $\tilde{t} = -0.1$. At the upper bounds of the Coulomb interaction we are in the insulator regime. In Fig. 3 the average density difference $\tilde{n}$ is presented along the excitonic pairing boundary, where the pair formation begins, thus at the transition from the symmetric $\Delta = 0$ to the unsymmetric phase $\Delta \neq 0$. We see that in this case we are starting nearly with the same values for the average densities n_a and n_b as in the case of the pair formation i.e. $n_a \approx 0.39$ and $n_b \approx 0.61$ at $U = 0$. They are different for the intermediate values of the Coulomb interaction parameter. It is clear from Fig. 3 that, approaching to the boundary at which $\Delta = 0$, we are reaching the insulator state more drastically than in the case with the excitonic pairs. Indeed, for example at $U/t = 6$ we have $n_a = 0.107$ and $n_b = 0.893$ at the boundary $\Delta = 0$, while, in the case when $\Delta \neq 0$, we have for the same value of the Coulomb interaction $n_a = 0.1625$ and $n_b = 0.8375$. Thus, approaching to the excitonic pairing boundary, the average density of the valence band electrons is slightly increasing. The lines in Fig. 4 represent the corresponding chemical potentials μ_Δ, along the excitonic transition boundary. The upper critical line corresponds to the case $\tilde{t} = -0.3$ and for the lower curve $\tilde{t} = -0.1$. The single particle excitation energies at the pair formation boundaries are sensible to the changes of the hopping amplitude $\tilde{t}$. Smaller values of $\tilde{t}$ correspond to the smaller excitation energies.

The chemical potential, in the case of the presence of excitonic pairs (derived from the system of equations in Eq.(61)), forms a well defined band and, consequently, implies the formation of the single particle excitation gap $\Delta_g = \mu^{max} - \mu^{min}$, where μ^{max} and μ^{min} are the upper and lower bounds of the chemical potential. The lower bounds of the chemical potentials for different values of the hopping integral $\tilde{t}$ are presented on the Fig. 5. We see from Fig. 5 that the evolution of the chemical potential is continuous from weak to strong coupling limit. It is worth to mention that the similar behavior for the chemical potential is found in Ref. 26, where the authors examine the Bose-condensation in an attractive interacting fermion gaz. In the inset of the Fig. 5 the average density difference $\tilde{n}$ is presented as a function of the excitonic gap parameter normalized to the hopping amplitude. For the small values of the Coulomb repulsion the single particle gap is totally degenerated and we have $\Delta_g = 0$. In this region the chemical potential is coinciding with the values presented on the Fig. 4. In a weak to intermediate coupling regime, the single particle gap Δ_g and the excitonic gap Δ both are increasing while in the strong couping limit ($U > 8$) the excitonic pairing gap parameter decreases rapidly with increasing U (see Fig. 2) and the single particle excitation gap remains open.



FIG. 3: The average particle density difference function $\tilde{n}$ along the pair formation transition boundary ($\Delta = 0$) as a function of the dimensionless interaction parameter U/t. Two different values of the hopping integral $\tilde{t}$ are considered.



FIG. 4: The chemical potentials along the pair formation transition boundary ($\Delta = 0$) as a function of the dimensionless interaction parameter U/t. Two different values of the hopping integral $\tilde{t}$ are considered. The chemical potentials are completely degenerated at the transition lines.



FIG. 5: The chemical potentials accompanying the excitonic pair formation as a function of the dimensionless interaction parameter U/t and at $T = 0$ K. Two different values of the hopping integral $\tilde{t}$ are considered. Degeneracy of the chemical potentials is lifted due to the EI state. Only the lower bounds of the chemical potentials are presented. In the inset, the variation of the density difference between two bands is presented as a function of the excitonic gap parameter for $\tilde{t} = -0.3$.

V. PHASE COHERENCE TRANSITION LINES

A. Phase stiffness at $T = 0$ K

In this section we examine the phase stiffness mechanism between the conduction band electrons and valence band holes. We will show how the phase coherence between these two solid state bands is important for the excitonic condensation order at the very low temperatures. For considering the phase stiffness we are interesting in purely phase action and we will integrate out the fermions in Eq.(32). Then we obtain the effective phase action. The partition function after integration is

$$\mathcal{Z} = \int \left[\mathcal{D}\varphi\right] e^{-\mathcal{S}_{\rm eff}[\varphi]}, \tag{68}$$

where

$$\mathcal{S}_{\rm eff}[\varphi] = -\ln \int \left[\mathcal{D}\bar{a}\mathcal{D}a\right] \left[\mathcal{D}\bar{b}\mathcal{D}b\right] e^{-\mathcal{S}}. \tag{69}$$

Basing on the self-consistency results presented in the Appendix A, we will write the effective phase action in the following form

$$\begin{aligned}\mathcal{S}_{\rm eff}[\varphi] = \;& \tilde{\mathcal{S}}_0 + \langle \mathcal{S}\rangle_{\mathcal{S}[\bar{a},a,\bar{b},b]} \\ & -\frac{1}{2}\left[\langle \mathcal{S}^2\rangle_{\mathcal{S}[\bar{a},a,\bar{b},b]} - \langle \mathcal{S}\rangle^2_{\mathcal{S}[\bar{a},a,\bar{b},b]}\right].\end{aligned} \tag{70}$$

The first term in Eq.(71) could be omitted in the calculations, because it gives the constant contribution to the total phase action. The averages in the right hand side of Eq.(70) are given in the sense that is discussed in Appendix A. From the second term in Eq.(70) only the first term in the Eq.(30) survives. Because we are searching the excitonic pairing and phase coherence mechanisms between the electrons and the holes in the system, we concentrate on the third term in Eq.(70) proportional to the average of the second order action and especially on the terms, which are proportional to the mixed products of the hopping amplitudes of two solid state bands $t\tilde{t}$ or $\tilde{t}t$. These terms are responsible for the phase stiffness. We can rewrite the action Eq.(70) in the form

$$\mathcal{S}_{\rm eff}[\varphi] = \tilde{\mathcal{S}}_0 + \mathcal{S}_0[\varphi] - \frac{1}{2}\left\langle \mathcal{S}^2\right\rangle_{\mathcal{S}[\bar{a},a,\bar{b},b]}, \tag{71}$$

where the action $\mathcal{S}_0[\varphi]$ is given in the Eq.(31). The detailed calculation of the average of the second order action in Eq.(71) is given in the Appendix B. Keeping in evaluations only the terms proportional to the second order in the hopping amplitudes, we have for the phase stiffness action

$$\mathcal{S}_J\left[\varphi\right] = -J\int_0^\beta d\tau \sum_{\langle \mathbf{r}\mathbf{r}'\rangle} \cos 2\left[\varphi(\mathbf{r}\tau) - \varphi(\mathbf{r}'\tau)\right], \tag{72}$$

where J is the phase stiffness parameter. The evaluations of J are given in details in the Appendix B and here we present only the phase stiffness critical lines. On the Fig. 6 we presented the phase stiffness parameter J for two different values of the hopping amplitude $\tilde{t}$. As the figures show, the phase stiffnesses (see the Appendix B) are strictly positive for all values of the Coulomb interaction. The upper curve in Fig. 6 corresponds to $\tilde{t} = -0.3$ and for the lower curve we have chosen the value $\tilde{t} = -0.1$. The phase stiffness is negligibly small, but is present along the whole EI region and it is especially important for the intermediate values of the Coulomb interaction, where the system of excitons undergoes the Bose-superfluid condensation.

B. The condensate order

Following discussion provided in the Section IV, the excitonic gap parameter at the very low temperatures (when the on-condensate excitations are negligible) could be written as in Eq.(60) with the help of the condensate transition probability amplitude ψ_0. In this section we will derive the excitonic condensate transition temperature in the frames of the quantum rotor model. In the discussion above, we derived the form of the effective φ-action $\mathcal{S}_{\rm eff}\left[\varphi\right] = \mathcal{S}_0\left[\varphi\right] + \mathcal{S}_J\left[\varphi\right]$. Here we start with the partition function given by



FIG. 6: The zero temperature phase stiffness parameter J given in Eq.(72) for the three-dimensional system as a function of the dimensionless interaction parameter U/t and for two different values of the hopping parameter $\tilde{t}$.

$$\mathcal{Z} = \int \left[\mathcal{D}\varphi\right] e^{-\mathcal{S}_{\rm eff}[\varphi]}, \tag{73}$$

To proceed, we replace the phase degrees of freedom by the complex unimodular field $z(\mathbf{r}\tau)$

$$z(\mathbf{r}\tau) = e^{i\varphi(\mathbf{r}\tau)}. \tag{74}$$

The variables $z(\mathbf{r}\tau)$ satisfy also the periodic boundary conditions $z(\mathbf{r}\beta) = z(\mathbf{r}0)$. This is done by implementing these new complex variables with the Faddeev-Popov ghost-fields method,[27] and we can introduce into the partition function in Eq.(73) the following identity

$$\begin{aligned}&\int \mathcal{D}\bar{z}\mathcal{D}z\delta\left(\sum_{\mathbf{r}}|z(\mathbf{r}\tau)|^2 - N\right)\times \\ &\times\delta(z - e^{i\varphi(\mathbf{r}\tau)})\delta(\bar{z} - e^{-i\varphi(\mathbf{r}\tau)}) = 1.\end{aligned} \tag{75}$$

We introduce then the spherical constraint for the unimodular complex variables $z(\mathbf{r}\tau)$ ($|z(\mathbf{r}\tau)|^2 = 1$). Indeed, the Eq.(75) implies the evident condition

$$\sum_{\mathbf{r}}|z(\mathbf{r}\tau)|^2 = N. \tag{76}$$

The constraint of unimodularity can be resolved by introducing the Lagrange multiplier λ and using the functional representation of the Dirac delta-function. This adds a quadratic term (in the z-field) to the phase action

$$\delta\left(\sum_{\mathbf{r}}|z(\mathbf{r}\tau)|^2-N\right)=\int_{-i\infty}^{+i\infty}\left[\frac{\mathcal{D}\lambda}{2\pi i}\right]\times$$
$$\times e^{-i\int_0^\beta d\tau\sum_{\mathbf{r}}\lambda\left(\,|z(\mathbf{r}\tau)|^2-1\,\right)}. \qquad (77)$$

Such a description is justified in the sense that the pair condensate has the macroscopic quantum phase coherence with the order parameter ψ_0 (see the section IV). The phase stiffness action in Eq.(72) can be rewritten in more convenient form using the trigonometric transformation formula

$$\cos 2\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau)\right]=2\cos^2\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau)\right]-1. \qquad (78)$$

Then, in terms of the complex variables $z(\mathbf{r}\tau)$, the transformation in Eq.(78) leads to a biquadratic form of the phase stiffness action. We have

$$\mathcal{S}_J\left[\bar{z},z\right]=-\frac{J}{2}\int_0^\beta d\tau\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}\left[\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau)+c.c.\right]^2. \qquad (79)$$

This biquadratic action could be transformed into a quadratic one with the help of the Hubbard-Stratanovich transformation for the one component bose field, by introducing the complex variables $w(\mathbf{r}\tau\mathbf{r}'\tau)$ at each pair of lattice positions $\mathbf{r}$, $\mathbf{r}'$ and time τ

$$e^{\frac{J}{2}\int_0^\beta d\tau\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}\left[\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau)+c.c.\right]^2}$$
$$=\int\left[\mathcal{D}w\right]e^{-\int_0^\beta d\tau\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}\frac{w(\mathbf{r}\tau\mathbf{r}'\tau)^2}{2J}}\times$$
$$\times e^{\int_0^\beta d\tau\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}w(\mathbf{r}\tau\mathbf{r}'\tau)\left[\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau)+c.c.\right]}. \qquad (80)$$

The saddle-point evaluation of the obtained action gives the critical value for the decoupling field $w(\mathbf{r}\tau\mathbf{r}'\tau)$

$$w_0=J\left\langle\left[\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau)+c.c.\right]\right\rangle\equiv 2Jg, \qquad (81)$$

where we used the expression of the bandwidth renormalization factor g given in the Eq.(42). Putting Eq.(81) back into the expression Eq.(80) we will have the phase stiffness contribution to the partition function in the spherical model

$$e^{4gJ\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}\int_0^\beta d\tau\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau)}. \qquad (82)$$

Hence, the effective phase stiffness action $\mathcal{S}_J[\bar{z},z]$, after the decoupling procedure, is reduced to a quadratic one in $z(\mathbf{r}\tau)$ field, i.e.

$$\mathcal{S}_J[\bar{z},z]=-4gJ\sum_{\langle\mathbf{r}\mathbf{r}'\rangle}\int_0^\beta d\tau\bar{z}(\mathbf{r}\tau)z(\mathbf{r}'\tau). \qquad (83)$$

Integrating the phase variables φ in Eq.(73), we will have for the partition function of our quantum rotor approach the following expression

$$\mathcal{Z}=\mathcal{Z}_0\int\left[\mathcal{D}\lambda\right]\int\left[\mathcal{D}\bar{z}\mathcal{D}z\right]e^{-\mathcal{S}_\lambda[\bar{z},z]}, \qquad (84)$$

where $\mathcal{Z}_0$ in Eq.(84) is given as

$$\mathcal{Z}_0=\int\left[\mathcal{D}\varphi\right]e^{-\mathcal{S}_0[\varphi]}. \qquad (85)$$

$\mathcal{S}_\lambda[\bar{z},z]$ in Eq.(84) is the total spherical action for our model. The form of the action $\mathcal{S}_\lambda[\bar{z},z]$ in the Fourier space is

$$\begin{aligned}\mathcal{S}_\lambda[\bar{z},z] &= \frac{1}{\beta N}\sum_{\mathbf{k}\omega_n}\bar{z}(\mathbf{k}\omega_n)\mathcal{G}_z^{-1}(\mathbf{k}\omega_n)z(\mathbf{k}\omega_n)\\ &-\beta N\lambda,\end{aligned} \qquad (86)$$

where

$$\mathcal{G}_z^{-1}(\mathbf{k}\omega_n)=\lambda-8gJ\epsilon(\mathbf{k})+\gamma^{-1}(\omega_n) \qquad (87)$$

is the inverse of the bosonic charge propagator and ω_n in Eq.(86) are the Bose-Matsubara frequencies. $\gamma^{-1}(\omega_n)$ is the inverse of the Fourier transformation of the two-point phase correlator $\gamma(\mathbf{r}\tau,\mathbf{r}'\tau')$ appearing after the phase integration. It is defined as

$$\gamma(\mathbf{r}\tau,\mathbf{r}'\tau')=\frac{1}{\mathcal{Z}_0}\int\left[\mathcal{D}\varphi\right]e^{-\mathcal{S}_0[\varphi]}e^{i\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}'\tau')\right]}. \qquad (88)$$

The calculation of the Fourier transformation $\gamma(\omega_n)$ of the function in Eq.(88) is straightforward[29]

$$\gamma(\omega_n)=\frac{8}{U\mathcal{Z}_0}\sum_{m=-\infty}^{+\infty}\frac{e^{-\frac{U\beta}{4}\left(m-\frac{\bar{\mu}}{U}\right)^2}}{1-16\left[\frac{i\omega_n}{U}-\left(m-\frac{\bar{\mu}}{U}\right)\right]^2}, \qquad (89)$$

where

$$\mathcal{Z}_0=\sum_{m=-\infty}^{+\infty}e^{-\frac{U\beta}{4}\left(m-\frac{\bar{\mu}}{U}\right)^2} \qquad (90)$$

and the summation is over the winding numbers m, which define a manifold composed of an infinite set of the topologically equivalent paths. Along the critical line the inverse of the uniform static order parameter susceptibility vanishes

$$\mathcal{G}_z^{-1}(\mathbf{k}=\mathbf{0},\omega_n=0)=0, \qquad (91)$$

which fixes the value of the Lagrange multiplier to the value $\lambda=\lambda_0$ given by the equation

$$\lambda_0-8gJ\epsilon(\mathbf{k}=0)+\gamma^{-1}(\omega_n=0)=0. \qquad (92)$$

At the condensate phase transition boundary, we have the equation for the critical temperature

$$\frac{1}{\beta N}\sum_{\mathbf{k}\omega_n}\frac{1}{\lambda_0-8gJ\epsilon(\mathbf{k})+\gamma^{-1}(\omega_n)}=1. \qquad (93)$$

Considering the principal $m = 0$ topological path, which is essential for the Bose-superfluid condensation, and performing the Bose-Matsubara frequency summation in Eq.(93), we will have the following self-consistent equation for the critical temperature

$$\frac{U}{2N}\sum_{\mathbf{k}}\frac{n_B\left(\zeta_{1\mathbf{k}}\right)-n_B\left(\zeta_{1\mathbf{k}}\right)}{\sqrt{\bar{\mu}^2+16UgJ\left[\epsilon(\mathbf{0})-\epsilon(\mathbf{k})\right]}}=1, \tag{94}$$

where $n_B\left(\epsilon\right)$ is the Bose-Einstein distribution function $n_B\left(\epsilon\right) = 1/\left(e^{\beta\epsilon}-1\right)$ and the variables $\zeta_{1\mathbf{k}}$ and $\zeta_{2\mathbf{k}}$ are introduced as

$$\zeta_{1\mathbf{k}} = \frac{1}{2}\left[-\bar{\mu}+\sqrt{\bar{\mu}^2+16UJg\left[\epsilon(\mathbf{0})-\epsilon(\mathbf{k})\right]}\right], \tag{95}$$

$$\zeta_{2\mathbf{k}} = \frac{1}{2}\left[-\bar{\mu}-\sqrt{\bar{\mu}^2+16UJg\left[\epsilon(\mathbf{0})-\epsilon(\mathbf{k})\right]}\right]. \tag{96}$$

$\epsilon(\mathbf{k})$ stands for the lattice dispersion for the three-dimensional cubic system and is given in Eq.(41). The simultaneous solution of the equation Eq.(94) with the system of self-consistent equations Eq.(61) from the Section IV gives the temperature transition critical line to the excitonic condensate state (EC) for the Falicov-Kimball model. The result is presented on the Fig. 7 (the lower curve on the figure). The upper curve in the panel represents the temperature transition to the excitonic pair state (EP) for $\tilde{t} = -0.3$. This transition line defines the phase boundary at which the excitonic pair formation starts. Above the transition line the system is in the normal Fermi Liquid[28] (FL) state (N). Passing from the strong to the weak interaction regime, we observe that the macroscopic condensate begins its formation practically at the same time with the excitonic pair formation and vanishes in a very weak interaction region (nearly at the same time that the exciton gap order disappears). For $\tilde{t} = -0.1$ we have presented the solution on the Fig. 8. We see that with decreasing the hopping parameter for the initial f-orbital electrons we obtain the similar temperature transition lines but with lower transition amplitudes, so diminishing the hopping energy has a destructive influence on the condensate transition and for a very low hopping amplitudes we could have the total destruction of the condensate state. From this point of observations, we can suggest that the condensate phase is due to the itinerant electronic correlations and localizing them leads to the destruction of the excitonic condensate state. It is clear that even very small phase correlations as exist in our model lead to the comparatively large (in amplitude) phase transition into the condensate regime with non-zero critical temperature. The maximum of the condensate transition coincides with the maximum of the phase stiffness plot (see Fig. 6) and this is due to the fact that both transitions are strongly related. From the behavior of the excitonic gap parameter (Fig. 2) it is clear that the transition maximum is displaced from that of the phase stiffness (Fig. 6) and the condensate order (Fig. 7). The maximum for the gap (in the case of $\tilde{t} = -0.3$) is achieved at $U/t = 7.8$, which corresponds to an imbalance of the c and f band electron concentrations $n_b - n_a = 0.8$ (this is near to the insulator regime), while the maximums for J and T_c are pointed nearly at the same value of $U/t = 5.1$, which corresponds to the limit of semiconductor with nearly equal values of the electron concentrations $n_a \approx n_b$. Physically, this is due also to the fact that the excitonic pair formation begins at the same time that the phase coherence appears in the system but the maximum number of excitonic pairs in the system is reached in the strong interaction region, while for the condensed pairs the maximum is shifted to the small and medium interaction regions. Another interesting physical interpretation of this phenomenon is that the phase coherence in the system is responsible for the Bose-superfluid phase transition and might cause also the excitonic pair formation, but, in essence, they still uncorrelated.



FIG. 7: The transition temperature lines in the three-dimensional Falicov-Kimball model for $\tilde{t} = -0.3$ as a function of the normalized Coulomb interaction strength U. The upper curve represents the excitonic pair (EP) transition temperature line T_Δ i.e. when $\Delta = 0$ and $\psi_0 = 0$. The lower curve represents the excitonic condensate (EC) transition temperature T_c with $\Delta \neq 0$ and $\psi_0 \neq 0$.

VI. SUMMARY

In the present paper we have shown the existence of the Bose-superfuild condensate order in the three-dimensional strongly correlated electron system. We considered the phase stiffness and we derived the effective phase stiffness action in the model. On the base of the



FIG. 8: The transition temperature lines in the three-dimensional Falicov-Kimball model for $\tilde{t} = -0.1$ as a function of the normalized Coulomb interaction strength U. The upper curve represents the excitonic pair (EP) transition temperature line T_Δ i.e. when $\Delta = 0$ and $\psi_0 = 0$. The lower curve represents the excitonic condensate (EC) transition temperature T_c with $\Delta \neq 0$ and $\psi_0 \neq 0$.

phase coherence between the conduction band electrons and valence band holes, we derived the expression for the phase stiffness parameter and we have shown that it is positive along the whole pair formation region of the normalized Coulomb interaction parameter. We proved that the phase stiffness is important for the macroscopic phase coherence order in the system, leading to the condensation of the preformed excitonic pairs into the fundamental state of the reciprocal momentum state. From our results it is clear that the critical temperature of the excitonic pair formation is not coinciding with the Bose-superfluid critical temperature. The Bose condensation of the excitonic pairs appears at the temperatures much smaller than the excitonic pair formation critical temperature is. Another important result is that of the existence of the BCS-BEC type crossover. In the small interaction limit the excitonic pairing and the condensate critical temperature lines are strongly coinciding. This is consistent with the superconducting transition scenario of the BCS model, in which the Cooper-pair formation and its condensation occur simultaneously and the system is in the BSC-like regime. For the strong interaction case we have the separation of the condensate order and the excitonic pairing critical temperature lines and the system is formally in the BEC-like regime. As a possible evaluation of our results in future would be the determination of the single-particle excitation spectra and density of states in the frames of the EFKM model. This will give us an access to compare our theoretical results with the temperature-dependent photoemission spectroscopy measurements.

Acknowledgments

Appendix A: The method of effective actions

1. Fermionic action

We start here with the partition function given in the Eq.(32). We treat our problem of strongly correlated electrons in a self-consistent-field approximation based on the U(1) gauge transformation introduced in the Section III. This is in general equivalent to the second order in the Fermi-liquid theory.[28] First, we integrate out the phase variables φ in Eq.(32)

$$\mathcal{Z} = \int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] e^{-\mathcal{S}[\bar{a},a,\bar{b},b]}, \tag{A1}$$

where the effective fermionic action $\mathcal{S}[\bar{a},a,\bar{b},b]$ in the exponential is defined as

$$\mathcal{S}[\bar{a},a,\bar{b},b] = -\ln \int \left[\mathcal{D}\varphi\right] e^{-\mathcal{S}[\bar{a},a,\bar{b},b,\varphi]}. \tag{A2}$$

Then we expand the logarithm keeping only the terms up to second order in the hopping amplitude. We get for the effective action

$$\begin{aligned}\mathcal{S}[\bar{a},a,\bar{b},b] = \quad & \mathcal{S}_0 + \langle\mathcal{S}\rangle_{\mathcal{S}_\varphi} \\ & -\frac{1}{2}\left[\left\langle\mathcal{S}^2\right\rangle_{\mathcal{S}_\varphi} - \langle\mathcal{S}\rangle^2_{\mathcal{S}_\varphi}\right].\end{aligned} \tag{A3}$$

Here

$$\langle\ldots\rangle_\varphi = \frac{\int\left[\mathcal{D}\varphi\right]\ldots e^{-\mathcal{S}}}{\int\left[\mathcal{D}\varphi\right]e^{-\mathcal{S}}}. \tag{A4}$$

The phase averaging can be considered as the quantum statistical averaging with an effective phase field action $\mathcal{S}_{\rm eff}[\varphi]$. This consideration is sometimes called like the self-consistency condition. Thus, we replace

$$\langle\mathcal{S}\rangle_\varphi \to \langle\mathcal{S}\rangle_{\mathcal{S}_{\rm eff}[\varphi]} \tag{A5}$$

and

$$\langle\ldots\rangle_{\mathcal{S}_{\rm eff}[\varphi]} = \frac{\int\left[\mathcal{D}\varphi\right]\ldots e^{-\mathcal{S}_{\rm eff}[\varphi]}}{\int\left[\mathcal{D}\varphi\right]e^{-\mathcal{S}_{\rm eff}[\varphi]}}. \tag{A6}$$

Then, for the effective fermionic action we have

$$\begin{aligned}\mathcal{S}[\bar{a},a,\bar{b},b] = \quad & \mathcal{S}_0 + \langle\mathcal{S}\rangle_{\mathcal{S}_{\rm eff}[\varphi]} \\ & -\frac{1}{2}\left[\left\langle\mathcal{S}^2\right\rangle_{\mathcal{S}_{\rm eff}[\varphi]} - \langle\mathcal{S}\rangle^2_{\mathcal{S}_{\rm eff}[\varphi]}\right].\end{aligned} \tag{A7}$$

2. Phase action

The integration over the fermions gives the effective action for the phase sector discussed above. The partition function in this case is

$$\mathcal{Z} = \int \left[\mathcal{D}\varphi\right] e^{-\mathcal{S}_{\text{eff}}[\varphi]}, \tag{A8}$$

where the effective phase action in the exponential is

$$\mathcal{S}_{\text{eff}}[\varphi] = -\ln \int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] e^{-\mathcal{S}}. \tag{A9}$$

Again, expanding the logarithm we will have

$$\begin{aligned}\mathcal{S}_{\text{eff}}[\varphi] = \quad & \tilde{\mathcal{S}}_0 + \left\langle \mathcal{S} \right\rangle_{ab} \\ & -\frac{1}{2}\left[\left\langle \mathcal{S}^2 \right\rangle_{ab} - \left\langle \mathcal{S} \right\rangle_{ab}^2\right].\end{aligned} \tag{A10}$$

Here the fermionic average $\left\langle ... \right\rangle_{ab}$ is

$$\left\langle ... \right\rangle_{ab} = \frac{\int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] ... e^{-\mathcal{S}}}{\int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] e^{-\mathcal{S}}}. \tag{A11}$$

Repeating the same steps as in the previous case and using the self-consistency condition

$$\left\langle \mathcal{S} \right\rangle_{ab} \to \left\langle \mathcal{S} \right\rangle_{\mathcal{S}\left[\bar{a},a,\bar{b},b\right]} \tag{A12}$$

we get for the effective phase action in this case

$$\begin{aligned}\mathcal{S}_{\text{eff}}[\varphi] = \quad & \tilde{\mathcal{S}}_0 + \left\langle \mathcal{S} \right\rangle_{\mathcal{S}\left[\bar{a},a,\bar{b},b\right]} \\ & -\frac{1}{2}\left[\left\langle \mathcal{S}^2 \right\rangle_{\mathcal{S}\left[\bar{a},a,\bar{b},b\right]} - \left\langle \mathcal{S} \right\rangle_{\mathcal{S}\left[\bar{a},a,\bar{b},b\right]}^2\right],\end{aligned} \tag{A13}$$

where $\tilde{\mathcal{S}}_0$ is an unimportant constant. After the self-consistency assumption, the average in Eq.(A11) will be

$$\left\langle ... \right\rangle_{\mathcal{S}\left[\bar{a},a,\bar{b},b\right]} = \frac{\int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] ... e^{-\mathcal{S}\left[\bar{a},a,\bar{b},b\right]}}{\int \left[\mathcal{D}\bar{a}\mathcal{D}a\right]\left[\mathcal{D}\bar{b}\mathcal{D}b\right] e^{-\mathcal{S}\left[\bar{a},a,\bar{b},b\right]}}. \tag{A14}$$

The relations in Eq.(A6) and Eq.(A14) are principal for the theory. Using these relations together, we resolve the problem of interacting fermions as a self-consistently coupled problem of the noninteracting fermions and the gauge-bosonic phase field. The bosonic sector plays the role of a suitable background, on which the collective excitations in the fermionic sector appear.

Appendix B: Phase stiffness

Here, we present a fermion averaging routine for averaging the four-fermionic type terms appearing in the expression of the third term of the effective phase action given in Eq.(71). This method is called the Wick averaging procedure and is used in many quantum-field theoretical treatments (see for example in Ref. 24). We present here in details the derivation of the term proportional to $t\tilde{t}$. The other term proportional to $\tilde{t}t$ can be derived analogically. For the fermion orbitals the mixed term proportional to $t\tilde{t}$ is

$$\begin{aligned}&-\frac{1}{2}\sum_{\left\langle \mathbf{r}_1\mathbf{r}'_1\right\rangle}\sum_{\left\langle \mathbf{r}_2\mathbf{r}'_2\right\rangle}\int_0^\beta d\tau d\tau' \left[t(\mathbf{r}_1\mathbf{r}'_1)\tilde{t}(\mathbf{r}_2\mathbf{r}'_2)\times\right. \\ &\left\langle \bar{a}(\mathbf{r}_1\tau)a(\mathbf{r}'_1\tau)\bar{b}(\mathbf{r}_2\tau')b(\mathbf{r}'_2\tau')\right\rangle \times \\ &\times e^{-i\left[\varphi(\mathbf{r}_1\tau)-\varphi(\mathbf{r}'_1\tau)\right]}e^{-i\left[\varphi(\mathbf{r}_2\tau')-\varphi(\mathbf{r}'_2\tau')\right]} \\ &+t(\mathbf{r}_1\mathbf{r}'_1)\tilde{t}(\mathbf{r}'_2\mathbf{r}_2)\left\langle \bar{a}(\mathbf{r}_1\tau)a(\mathbf{r}'_1\tau)\bar{b}(\mathbf{r}'_2\tau')b(\mathbf{r}_2\tau')\right\rangle \times \\ &\times e^{-i\left[\varphi(\mathbf{r}_1\tau)-\varphi(\mathbf{r}'_1\tau)\right]}e^{i\left[\varphi(\mathbf{r}_2\tau')-\varphi(\mathbf{r}'_2\tau')\right]} \\ &+t(\mathbf{r}'_1\mathbf{r}_1)\tilde{t}(\mathbf{r}_2\mathbf{r}'_2)\left\langle \bar{a}(\mathbf{r}'_1\tau)a(\mathbf{r}_1\tau)\bar{b}(\mathbf{r}_2\tau')b(\mathbf{r}'_2\tau')\right\rangle \times \\ &\times e^{i\left[\varphi(\mathbf{r}_1\tau)-\varphi(\mathbf{r}'_1\tau)\right]}e^{-i\left[\varphi(\mathbf{r}_2\tau')-\varphi(\mathbf{r}'_2\tau')\right]} \\ &+t(\mathbf{r}'_1\mathbf{r}_1)\tilde{t}(\mathbf{r}'_2\mathbf{r}_2)\left\langle \bar{a}(\mathbf{r}'_1\tau)a(\mathbf{r}_1\tau)\bar{b}(\mathbf{r}'_2\tau')b(\mathbf{r}_2\tau')\right\rangle \times \\ &\left.\times e^{i\left[\varphi(\mathbf{r}_1\tau)-\varphi(\mathbf{r}'_1\tau)\right]}e^{i\left[\varphi(\mathbf{r}_2\tau')-\varphi(\mathbf{r}'_2\tau')\right]}\right].\end{aligned} \tag{B1}$$

As an example, we give the Wick averaging result of the first four-fermion term in the expression of Eq.(B1)

$$\begin{aligned}&\left\langle \bar{a}(\mathbf{r}_1\tau)a(\mathbf{r}'_1\tau)\bar{b}(\mathbf{r}_2\tau')b(\mathbf{r}'_2\tau')\right\rangle = \\ &\left\langle \bar{a}(\mathbf{r}_1\tau)a(\mathbf{r}'_1\tau)\right\rangle\left\langle \bar{b}(\mathbf{r}_2\tau')b(\mathbf{r}'_2\tau')\right\rangle \\ &-\left\langle \bar{a}(\mathbf{r}_1\tau)\bar{b}(\mathbf{r}_2\tau')\right\rangle\left\langle a(\mathbf{r}'_1\tau)b(\mathbf{r}'_2\tau')\right\rangle \\ &\left\langle \bar{a}(\mathbf{r}_1\tau)b(\mathbf{r}'_2\tau')\right\rangle\left\langle a(\mathbf{r}'_1\tau)\bar{b}(\mathbf{r}_2\tau')\right\rangle \\ &= \mathcal{G}^{aa}(\mathbf{r}'_1-\mathbf{r}_1,0)\mathcal{G}^{bb}(\mathbf{r}'_2-\mathbf{r}_2,0) \\ &-\mathcal{G}^{ab}(\mathbf{r}_1-\mathbf{r}'_2,\tau-\tau')\mathcal{G}^{ba}(\mathbf{r}_2-\mathbf{r}'_1,\tau'-\tau).\end{aligned} \tag{B2}$$

We will keep only the terms proportional to the electron density type order and the excitonic gap order. We neglect the other term containing the superconducting type ordering $\left\langle \bar{a}(\mathbf{r}\tau)\bar{b}(\mathbf{r}'\tau')\right\rangle$ or $\left\langle a(\mathbf{r}\tau)b(\mathbf{r}'\tau')\right\rangle$ because of the symmetry of the action in Eq.(36). In turn, for the first treatment, the terms containing the density type orderings $\left\langle \bar{a}(\mathbf{r}\tau)a(\mathbf{r}'\tau')\right\rangle$ and $\left\langle \bar{b}(\mathbf{r}\tau)b(\mathbf{r}'\tau')\right\rangle$ could be also omitted, since they are not contributing to the excitonic pair formation and condensate order. After calculating all averages in Eq.(B1) and recombining them with the similar terms coming from the component proportional to $\tilde{t}t$, we obtain the phase stiffness action in the form

$$\mathcal{S}_J\left[\varphi\right] = \int_0^\beta d\tau \sum_{\mathbf{r}\mathbf{r}'} \left\{ J(\mathbf{r}\tau,\mathbf{r}'\tau')\cos\left[\varphi(\mathbf{r}\tau)+\varphi(\mathbf{r}\tau')-\varphi(\mathbf{r}'\tau)-\varphi(\mathbf{r}'\tau')\right] \right.$$
$$\left. +\mathcal{G}_{ab}(\mathbf{0},\tau-\tau')\mathcal{G}_{ba}(\mathbf{0},\tau'-\tau)\cos\left[\varphi(\mathbf{r}\tau)-\varphi(\mathbf{r}\tau')-\varphi(\mathbf{r}'\tau)+\varphi(\mathbf{r}'\tau')\right]\right\}, \tag{B3}$$

where we introduced the interaction parameter J as

$$J = -4t\tilde{t}\int_0^\beta d\tau'\mathcal{G}_{ab}(\mathbf{r}-\mathbf{r}',\tau-\tau')\mathcal{G}_{ba}(\mathbf{r}-\mathbf{r}',\tau'-\tau). \tag{B4}$$

The fermion propagators in Eq.(B4) are defined in the Section IV. The second term in Eq.(B3) gives the constant contribution to the phase stiffness action. The variation of the product of the anomalous propagators with time difference $|\tau-\tau'|$ in the first term in Eq.(B3) is much faster than in the arguments of the trigonometric functions multiplied to it. For this reason, we will put $\tau'=\tau$ in all arguments of the trigonometric functions corresponding to the zero order term in the Taylor series expansion of the phase variables

$$\varphi(\mathbf{r}\tau') = \varphi(\mathbf{r}\tau) + (\tau'-\tau)\,\partial_\tau\varphi(\mathbf{r}\tau) + O\left[(\tau'-\tau)^2\right]. \tag{B5}$$

For the product of the anomalous propagators we have

$$\mathcal{G}_{ab}(\tau-\tau')\mathcal{G}_{ba}(\tau'-\tau) = \frac{4\Delta^2U^2}{z^2(\beta N)^2}\sum_{\mathbf{k}\mathbf{k}'}\sum_{\nu_n\nu'_n}\frac{\epsilon\left(\mathbf{k}\right)\epsilon\left(\mathbf{k}'\right)e^{-i(\nu'_n-\nu_n)\delta}}{\left[\mathcal{E}^a_{\mathbf{k}}(\nu_n)\mathcal{E}^b_{\mathbf{k}}(\nu_n)-|\Delta|^2\right]\left[\mathcal{E}^a_{\mathbf{k}'}(\nu'_n)\mathcal{E}^b_{\mathbf{k}'}(\nu'_n)-|\Delta|^2\right]}. \tag{B6}$$

Here $z=6$ is the number of the nearest neighbor sites on the three-dimensional cubic lattice. After integrating over the imaginary time in Eq.(B4), we perform the Matsubara summations over the fermionic Matsubara frequencies and we will have from the Eq.(B4) and Eq.(B6) the phase stiffness parameter

$$J = \frac{16\Delta^2t\tilde{t}}{z^2N^2}\sum_{\mathbf{k}\mathbf{k}'}\frac{\epsilon\left(\mathbf{k}\right)\epsilon\left(\mathbf{k}'\right)}{\sqrt{\xi^2_{\mathbf{k}}+4\Delta^2}}\left[\Lambda_1(\mathbf{k}\mathbf{k}')\tanh\left(\frac{\beta E^+_{\mathbf{k}}}{2}\right)-\Lambda_2(\mathbf{k}\mathbf{k}')\tanh\left(\frac{\beta E^-_{\mathbf{k}}}{2}\right)\right]. \tag{B7}$$

The parameters $\Lambda_1(\mathbf{k}\mathbf{k}')$ and $\Lambda_2(\mathbf{k}\mathbf{k}')$ entering in Eq.(B7) are given by

$$\Lambda_1(\mathbf{k}\mathbf{k}') = \frac{1}{E^+(\mathbf{k})-E^+(\mathbf{k}')}\cdot\frac{1}{E^+(\mathbf{k})-E^-(\mathbf{k}')},$$
$$\Lambda_2(\mathbf{k}\mathbf{k}') = \frac{1}{E^-(\mathbf{k})-E^-(\mathbf{k}')}\cdot\frac{1}{E^-(\mathbf{k})-E^+(\mathbf{k}')}. \tag{B8}$$

Following the Section IV, the summations over the $\mathbf{k}$-wave vectors in Eq.(B7) could be transformed into the integrations with the help of the appropriate density of states given in Eq.(67). The numerical evaluation results of the expression Eq.(B7) for $T=0$ K are presented in the Fig. 6 and discussed in the Section V of the present paper.

---

[1] Moskalenko S. A., Snoke D. W., "*Bose-Einstein Condensation of Excitons and Biexcitons*" (Cambridge University Press, 2005).

[2] L. V. Keldysh and A. N. Kozlov, Zh. Eksp. Teor. Fiz. **54**, 978993 (1968), Sov. Phys. JETP **27**, 521528 (1968).

[3] J. Neuenschwander, P. Wachter, Phys. Rev. B **41** (1990) 12693.

[4] P. Wachter, B. Bucher, J. Malar, Phys. Rev. B **69** (2004)

094502.

[5] L. V. Keldysh and Y. V. Kopaev, Soviet Phys. Solid State **6**, 2219 (1964).

[6] J. des Cloizeaux, J. Chem. Phys. Solids. 26, 259 (1965).

[7] W. Kohn, in Many Body Physics, edited by C. de Witt and R. Balian (Gordon & Breach, New York, 1968).

[8] D. Jérome, T. M. Rice, and W. Kohn, Phys. Rev. **158**, 462 (1967).

[9] L.V. Keldysh, in Bose-Einstein Condensation, A. Griffin, D.W. Snoke and S. Stringari, eds. (Cambridge U. Press, Cambridge, 1995).

[10] A. N. Kozlov, L. A. Maksimov, Sov. Phys. JETP, **21**, 790 (1965); **22**, 889 (1966).

[11] Y. Wakisaka, T. Sudayama, K. Takubo, T. Mizokawa, M. Arita, H. Namatame, M. Taniguchi, N. Katayama, M. Nohara, and H. Takagi, Phys. Rev. Lett. **103**, 026402 (2009).

[12] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **106**, 162, (1957).

[13] C. J. Pethick and H. Smith, "*BoseEinstein Condensation in Dilute Gases*", (Cambridge University Press, Cambridge, 2001).

[14] Qijin Chen, Jelena Stajic, Shina Tan, Kathryn Levin, Physics Reports **412**, 1-88 (2005).

[15] D. Kremp, D. Semkat, and K. Henneberger, Phys. Rev. B **78**, 125315 2008.

[16] B. Zenker, D. Ihle, F. X. Bronold, and H. Fehske, Phys. Rev. B **81**, 115122 March (2010).

[17] K. Seki, R. Eder, and Y. Ohta, Phys. Rev. B, **84**, 245106 (2011).

[18] B. Zenker, D. Ihle, F. X. Bronold, and H. Fehske, Phys. Rev. B **85**, 121102 (2012).

[19] L. M. Falicov and J. C. Kimball, Phys. Rev. Lett. **22**, 997 (1969).

[20] R. Ramirez, L. M. Falicov, and J. C. Kimball, Phys. Rev. B, **2**, 3383 (1970).

[21] E. Fradkin, "*Field Theories of Condensed Matter Systems*", (Addison- Wesley Publishing Company, 1991).

[22] L. Debski, Acta Physica Polonica A, **115,** No 1 (2009).

[23] A. V. Svidzinskii, "*Spatially Inhomogeneous Problems of Superconductivity Theory*", [in Russian], Nauka, Moscow (1982).

[24] G. D. Mahan, "*Many-particle physics*", (New York: Plenum, 1990).

[25] M. Abramovitz and I. Stegun, Handbook of Mathematical Functions (Dover, New York, 1970).

[26] P. Nozieres, S. Schmitt-Rink, Journal of Low Temperature Physics, **59**, Nos. 3/4, (1985).

[27] V. N. Popov, "*Functional Integrals in Quantum Field Theory and Statistical Physics*", (Springer, 2001).

[28] Ph. Nozieres and D. Pines, "*Theory Of Quantum Liquids*", (Westview Press, 1999).

[29] V. A. Apinyan and T. K. Kopeć, Phys. Rev. B, **78**, 184511 (2008).